

POSSIBLE RISKS, HAZARDS or COMPLICATIONS

(Please READ and INITIAL)

PAIN: There can be pain even after the topical anesthetic has been used. Anesthetics work better on some people than on others.
INFECTION: Infection is very unusual. The areas treated must be kept clean and only freshly cleaned hands should touch the areas. See "After Care" Instructions.
UNEVEN PIGMENTATION: This can result from poor healing, infection, bleeding, or many other causes. Your follow-up appointment will likely correct any uneven appearance.
ASYMMETRY: Every effort will be made to avoid asymmetry, but our faces are not symmetrical, so adjustments may be needed during the follow-up session to correct any unevenness.
EXCESSIVE SWELLING OR BRUISING: Some people bruise and swell more than others. Ice packs may help and the bruising and swelling typically disappears within 1-5 days. Most people don't bruise or swell at all.
ANESTHESIA: Topical anesthetics are used to numb the area to be tattooed. Lidocaine, Prilocaine, Benzocaine, Tetracain, and Epinephrine in a liquid, cream or gel forms are typically used. If you are allergic to any of these, please inform me immediately before procedure begins.
MRI: Because pigments used in permanent cosmetic procedures contain inert oxides, a low-level magnet may be required if you need to be scanned by an MRI machine. You must inform your technician of any tattoos or permanent cosmetics.
ALLERGIC REACTION
There is a small possibility of an allergic reaction. You may take a 5-7 day patch test to determine this.
Please initial to: A) WAIVE patch test B) TAKE Patch Test
The alternative to these possibilities is to use cosmetics and not undergo the Microblading / Microshading Procedure.
Consent and Release for procedures performed:
Signed Date